EGMO 2016, Day 1 – Solutions

Problem 1. Let n be an odd positive integer, and let x_1, x_2, ..., x_n be non-negative real numbers. Show that

$$\min_{i=1,\ldots,n}(x_i^2 + x_{i+1}^2) \leq \max_{j=1,\ldots,n}(2x_j x_{j+1}),$$

where $x_{n+1} = x_1$.

Solution. In what follows, indices are reduced modulo n. Consider the n differences $x_{k+1} - x_k$, $k = 1, \ldots, n$. Since n is odd, there exists an index j such that $(x_{j+1} - x_j)(x_{j+2} - x_{j+1}) \geq 0$. Without loss of generality, we may and will assume both factors non-negative, so $x_j \leq x_{j+1} \leq x_{j+2}$. Consequently,

$$\min_{k=1,\ldots,n}(x_k^2 + x_{k+1}^2) \leq x_j^2 + x_{j+1}^2 \leq 2x_{j+1}^2 \leq 2x_{j+1}x_{j+2} \leq \max_{k=1,\ldots,n}(2x_k x_{k+1}).$$

Remark. If $n \geq 3$ is odd, and one of the x_k is negative, then the conclusion may no longer hold. This is the case if, for instance, $x_1 = -b$, and $x_{2k} = a$, $x_{2k+1} = b$, $k = 1, \ldots, (n-1)/2$, where $0 \leq a < b$, so the string of numbers is

$$-b, b, a, b, a, \ldots, b, a.$$

If n is even, the conclusion may again no longer hold, as shown by any string of alternate real numbers: $a, b, a, b, \ldots, a, b$, where $a \neq b$.

Problem 2. Let $ABCD$ be a cyclic quadrilateral, and let diagonals AC and BD intersect at X. Let C_1, D_1 and M be the midpoints of segments CX, DX and CD, respectively. Lines AD_1 and BC_1 intersect at Y, and line MY intersects diagonals AC and BD at different points E and F, respectively. Prove that line XY is tangent to the circle through E, F and X.



Solution. We are to prove that $\backslash\, EXY = \backslash\, EFX$; alternatively, but equivalently, $\backslash\, AYX + \backslash\, XAY = \backslash\, BYF + \backslash\, XBY$.

Since the quadrangle $ABCD$ is cyclic, the triangles XAD and XBC are similar, and since AD_1 and BC_1 are corresponding medians in these triangles, it follows that $\backslash\, XAY = \backslash\, XAD_1 = \backslash\, XBC_1 = \backslash\, XBY$.

Finally, $\backslash\, AYX = \backslash\, BYF$, since X and M are corresponding points in the similar triangles ABY and C_1D_1Y: indeed, $\backslash\, XAB = \backslash\, XDC = \backslash\, MC_1D_1$, and $\backslash\, XBA = \backslash\, XCD = \backslash\, MD_1C_1$.

Problem 3. Let m be a positive integer. Consider a $4m \times 4m$ array of square unit cells. Two different cells are related to each other if they are in either the same row or in the same column. No cell is related to itself. Some cells are coloured blue, such that every cell is related to at least two blue cells. Determine the minimum number of blue cells.

Solution 1 (Israel). The required minimum is $6m$ and is achieved by a diagonal string of m 4×4 blocks of the form below (bullets mark centres of blue cells):

$$\begin{array}{cccc} \bullet & & & \\ \bullet & & & \\ \bullet & & & \\ & \bullet & \bullet & \bullet \end{array}$$

In particular, this configuration shows that the required minimum does not exceed $6m$.

We now show that any configuration of blue cells satisfying the condition in the statement has cardinality at least $6m$.

Fix such a configuration and let m_1^r be the number of blue cells in rows containing exactly one such, let m_2^r be the number of blue cells in rows containing exactly two such, and let m_3^r be the number of blue cells in rows containing at least three such; the numbers m_1^c, m_2^c and m_3^c are defined similarly.

Begin by noticing that $m_3^c \geq m_1^r$ and, similarly, $m_3^r \geq m_1^c$. Indeed, if a blue cell is alone in its row, respectively column, then there are at least two other blue cells in its column, respectively row, and the claim follows.

Suppose now, if possible, the total number of blue cells is less than $6m$. We will show that $m_1^r > m_3^r$ and $m_1^c > m_3^c$, and reach a contradiction by the preceding: $m_1^r > m_3^r \geq m_1^c > m_3^c \geq m_1^r$.

We prove the first inequality; the other one is dealt with similarly. To this end, notice that there are no empty rows — otherwise, each column would contain at least two blue cells, whence a total of at least $8m > 6m$ blue cells, which is a contradiction. Next, count rows to get $m_1^r + m_2^r/2 + m_3^r/3 \geq 4m$, and count blue cells to get $m_1^r + m_2^r + m_3^r < 6m$. Subtraction of the latter from the former multiplied by $3/2$ yields $m_1^r - m_3^r > m_2^r/2 \geq 0$, and the conclusion follows.

Solution 2. To prove that a minimal configuration of blue cells satisfying the condition in the statement has cardinality at least $6m$, consider a bipartite graph whose vertex parts are the rows and the columns of the array, respectively, a row and a column being joined by an edge if and only if the two cross at a blue cell. Clearly, the number of blue cells is equal to the number of edges of this graph, and the relationship condition in the statement reads: for every row r and every column c, $\deg r + \deg c - \epsilon(r, c) \geq 2$, where $\epsilon(r, c) = 2$ if r and c are joined by an edge, and $\epsilon(r, c) = 0$ otherwise.

Notice that there are no empty rows/columns, so the graph has no isolated vertices. By the preceding, the cardinality of every connected component of the graph is at least 4, so there are at most $2 \cdot 4m/4 = 2m$ such and, consequently, the graph has at least $8m - 2m = 6m$ edges. This completes the proof.

Remarks. The argument in the first solution shows that equality to $6m$ is possible only if $m_1^r = m_3^r = m_1^c = m_3^c = 3m$, $m_2^r = m_2^c = 0$, and there are no rows, respectively columns, containing four blue cells or more.

Consider the same problem for an $n \times n$ array. The argument in the second solution shows that the corresponding minimum is $3n/2$ if n is divisible by 4, and $3n/2 + 1/2$ if n is odd; if $n \equiv 2 \pmod 4$, the minimum in question is $3n/2 + 1$. To describe corresponding minimal configurations C_n, refer to the minimal configurations C_2, C_3, C_4, C_5 below:



The case $n \equiv 0 \pmod 4$ was dealt with above: a C_n consists of a diagonal string of $n/4$ blocks C_4. If $n \equiv r \pmod 4$, $r = 2, 3$, a C_n consists of a diagonal string of $\lfloor n/4 \rfloor$ blocks C_4 followed by a C_r, and if $n \equiv 1 \pmod 4$, a C_n consists of a diagonal string of $\lfloor n/4 \rfloor - 1$ blocks C_4 followed by a C_5.

Minimal configurations are not necessarily unique (two configurations being equivalent if one is obtained from the other by permuting the rows and/or the columns). For instance, if $n = 6$, the configurations below are both minimal:



EGMO 2016, Day 2 – Solutions

Problem 4. Two circles, ω_1 and ω_2, of equal radius intersect at different points X_1 and X_2. Consider a circle ω externally tangent to ω_1 at a point T_1, and internally tangent to ω_2 at a point T_2. Prove that lines X_1T_1 and X_2T_2 intersect at a point lying on ω.

Solution 1. Let the line X_kT_k and ω meet again at X_k', $k = 1, 2$, and notice that the tangent t_k to ω_k at X_k and the tangent t_k' to ω at X_k' are parallel. Since the ω_k have equal radii, the t_k are parallel, so the t_k' are parallel, and consequently the points X_1' and X_2' coincide (they are not antipodal, since they both lie on the same side of the line T_1T_2). The conclusion follows.

Solution 2. The circle ω is the image of ω_k under a homothety h_k centred at T_k, $k = 1, 2$. The tangent to ω at $X_k' = h_k(X_k)$ is therefore parallel to the tangent t_k to ω_k at X_k. Since the ω_k have equal radii, the t_k are parallel, so $X_1' = X_2'$; and since the points X_k, T_k and X_k' are collinear, the conclusion follows.





Solution 3. Invert from X_1 and use an asterisk to denote images under this inversion. Notice that ω_k^* is the tangent from X_2^* to ω^* at T_k^*, and the pole X_1 lies on the bisectrix of the angle formed by the ω_k^*, not containing ω^*. Letting $X_1T_1^*$ and ω^* meet again at Y, standard angle chase shows that Y lies on the circle $X_1^*X_2^*T_2^*$, and the conclusion follows.

Remarks. The product h_1h_2 of the two homotheties in the first solution is reflexion across the midpoint of the segment X_1X_2, which lies on the line T_1T_2.

Various arguments, involving similarities, radical axes, and the like, work equally well to prove the required result.

Problem 5. Let k and n be integers such that $k \geq 2$ and $k \leq n \leq 2k - 1$. Place rectangular tiles, each of size $1 \times k$ or $k \times 1$, on an $n \times n$ chessboard so that each tile covers exactly k cells, and no two tiles overlap. Do this until no further tile can be placed in this way. For each such k and n, determine the minimum number of tiles that such an arrangement may contain.

Solution. The required minimum is n if $n = k$, and it is $\min(n; 2n \quad 2k + 2)$ if $k < n < 2k$.

The case $n = k$ being clear, assume henceforth $k < n < 2k$. Begin by describing maximal arrangements on the board $[0; n] \quad [0; n]$, having the above mentioned cardinalities.

If $k < n < 2k \quad 1$, then $\min(n; 2n \quad 2k + 2) = 2n \quad 2k + 2$. To obtain a maximal arrangement of this cardinality, place four tiles, $[0; k] \quad [0; 1]$, $[0; 1] \quad [1; k + 1]$, $[1; k + 1] \quad [k; k + 1]$ and $[k; k + 1] \quad [0; k]$ in the square $[0; k] \quad [0; k]$, stack $n \quad k \quad 1$ horizontal tiles in the rectangle $[1; k + 1] \quad [k + 1; n]$, and erect $n \quad k \quad 1$ vertical tiles in the rectangle $[k + 1; n] \quad [1; k + 1]$.

If $n = 2k \quad 1$, then $\min(n; 2n \quad 2k + 2) = n = 2k \quad 1$. A maximal arrangement of $2k \quad 1$ tiles is obtained by stacking $k \quad 1$ horizontal tiles in the rectangle $[0; k] \quad [0; k \quad 1]$, another $k \quad 1$ horizontal tiles in the rectangle $[0; k] \quad [k; 2k \quad 1]$, and adding the horizontal tile $[k \quad 1; 2k \quad 1] \quad [k \quad 1; k]$.

The above examples show that the required minimum does not exceed the mentioned values.

To prove the reverse inequality, consider a maximal arrangement and let r, respectively c, be the number of rows, respectively columns, not containing a tile.

If $r = 0$ or $c = 0$, the arrangement clearly contains at least n tiles.

If r and c are both positive, we show that the arrangement contains at least $2n \quad 2k + 2$ tiles. To this end, we will prove that the rows, respectively columns, not containing a tile are consecutive. Assume this for the moment, to notice that these $r2$

to notice that these $r2$
to notice that these $r2$
to notice that these $r2$
to notice that th